

A COMPANION OF OSTROWSKI LIKE INEQUALITY FOR MAPPINGS WHOSE SECOND DERIVATIVES BELONG TO L^∞ SPACES AND APPLICATIONS

WENJUN LIU

ABSTRACT. A companion of Ostrowski like inequality for mappings whose second derivatives belong to L^∞ spaces is established. Applications to composite quadrature rules, and to probability density functions are also given.

1. INTRODUCTION

In 1938, Ostrowski established the following interesting integral inequality (see [15]) for differentiable mappings with bounded derivatives:

Theorem 1.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) whose derivative is bounded on (a, b) and denote $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$. Then for all $x \in [a, b]$ we have*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty.$$

The constant $\frac{1}{4}$ is sharp in the sense that it can not be replaced by a smaller one.

Motivated by [12], Dragomir [8] proved some companions of Ostrowski's inequality, as follows:

Theorem 1.2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$. Then the following inequalities*

$$(1.2) \quad \begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \begin{cases} \left[\frac{1}{8} + 2 \left(\frac{x - \frac{3a+b}{4}}{b-a} \right)^2 \right] (b-a) \|f'\|_\infty, & f' \in L^\infty[a, b], \\ \frac{2^{1/q}}{(q+1)^{1/q}} \left[\left(\frac{x-a}{b-a} \right)^{q+1} + \left(\frac{\frac{a+b}{2}-x}{b-a} \right)^{q+1} \right]^{1/q} (b-a)^{1/q} \|f'\|_p, & p > 1, \frac{1}{p} + \frac{1}{q} = 1 \text{ and } f' \in L^p[a, b], \\ \left[\frac{1}{4} + \left| \frac{x - \frac{3a+b}{4}}{b-a} \right| \right] \|f'\|_1, & f' \in L^1[a, b] \end{cases} \end{aligned}$$

hold for all $x \in [a, \frac{a+b}{2}]$.

Recently, Alomari [1] introduced a companion of Dragomir's generalization of Ostrowski's inequality for absolutely continuous mappings whose first derivatives are belong to $L^\infty([a, b])$.

2010 *Mathematics Subject Classification.* 26D15, 41A55, 41A80, 65C50.

Key words and phrases. Ostrowski like inequality; twice differentiable mapping; L^∞ spaces; composite quadrature rule; probability density function.

Theorem 1.3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous mappings on (a, b) whose derivative is bounded on $[a, b]$. Then the inequality*

$$(1.3) \quad \left| \left[(1-\lambda) \frac{f(x) + f(a+b-x)}{2} + \lambda \frac{f(a) + f(b)}{2} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \left[\frac{1}{8} (2\lambda^2 + (1-\lambda)^2) + 2 \frac{\left(x - \frac{(3-\lambda)a + (1+\lambda)b}{4} \right)^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

holds for all $\lambda \in [0, 1]$ and $x \in [a + \lambda \frac{b-a}{2}, \frac{a+b}{2}]$.

In (1.3), choose $\lambda = \frac{1}{2}$, one can get

$$(1.4) \quad \left| \frac{1}{2} \left[\frac{f(x) + f(a+b-x)}{2} + \frac{f(a) + f(b)}{2} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \left[\frac{3}{32} + 2 \frac{\left(x - \frac{5a+3b}{8} \right)^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty.$$

And if choose $x = \frac{a+b}{2}$, then one has

$$(1.5) \quad \left| \frac{1}{2} \left[f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8} (b-a) \|f'\|_\infty.$$

It's shown in [1] that the constant $\frac{1}{8}$ is the best possible.

In related work, Dragomir and Sofo [10] developed the following Ostrowski like integral inequality for twice differentiable mapping.

Theorem 1.4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a mapping whose first derivative is absolutely continuous on $[a, b]$ and assume that the second derivative $f'' \in L^\infty([a, b])$. Then we have the inequality*

$$(1.6) \quad \left| \frac{1}{2} \left[f(x) + \frac{f(a) + f(b)}{2} \right] - \frac{1}{2} \left(x - \frac{a+b}{2} \right) f'(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \left[\frac{1}{48} + \frac{1}{3} \frac{|x - \frac{a+b}{2}|^3}{(b-a)^3} \right] (b-a)^2 \|f''\|_\infty,$$

for all $x \in [a, b]$.

In (1.6), the authors pointed out that the midpoint $x = \frac{a+b}{2}$ gives the best estimator, i.e.,

$$(1.7) \quad \left| \frac{1}{2} \left[f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{48} (b-a)^2 \|f''\|_\infty.$$

In fact, we can choose $f(t) = (t-a)^2$ in (1.7) to prove that the constant $\frac{1}{48}$ in inequality (1.7) is sharp.

For other related results, the reader may be refer to [2, 3, 4, 5, 7, 9, 11, 13, 14, 16, 17, 18, 19, 20, 21] and the references therein.

Motivated by previous works [1, 6, 8, 10], we investigate in this paper a companion of the above mentioned Ostrowski like integral inequality for twice differentiable mappings. Our result gives a smaller estimator than (1.7) (see (2.9) below). Some applications to composite quadrature rules, and to probability density functions are also given.

2. A COMPANION OF OSTROWSKI LIKE INEQUALITY

The following companion of Ostrowski like inequality holds:

Theorem 2.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a mapping whose first derivative is absolutely continuous on $[a, b]$ and assume that the second derivative $f'' \in L^\infty([a, b])$. Then we have the inequality*

$$(2.1) \quad \left| \frac{1}{2} \left[\frac{f(x) + f(a+b-x)}{2} + \frac{f(a) + f(b)}{2} \right] - \frac{1}{2} \left(x - \frac{a+b}{2} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{3} \frac{(\frac{a+3b}{4} - x)(x-a)^2}{(b-a)^3} + \frac{1}{3} \frac{(\frac{a+b}{2} - x)^3}{(b-a)^3} \right] (b-a)^2 \|f''\|_\infty$$

for all $x \in [a, \frac{a+b}{2}]$. The first constant $\frac{1}{3}$ in the right hand side of (2.1) is sharp in the sense that it can not be replaced by a smaller one provided that $x \neq \frac{a+3b}{4}$ and $x \neq a$.

Proof. Define the kernel $K(t) : [a, b] \rightarrow \mathbb{R}$ by

$$(2.2) \quad K(t) := \begin{cases} t-a, & t \in [a, x], \\ t - \frac{a+b}{2}, & t \in (x, a+b-x], \\ t-b, & t \in (a+b-x, b], \end{cases}$$

for all $x \in [a, \frac{a+b}{2}]$. Integrating by parts, we obtain (see [8])

$$(2.3) \quad \frac{1}{b-a} \int_a^b K(t) g'(t) dt = \frac{g(x) + g(a+b-x)}{2} - \frac{1}{b-a} \int_a^b g(t) dt.$$

Now choose in (2.3), $g(x) = (x - \frac{a+b}{2})f'(x)$, to get

$$(2.4) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b K(t) \left[f'(t) + \left(t - \frac{a+b}{2} \right) f''(t) \right] dt \\ &= \frac{1}{2} \left(x - \frac{a+b}{2} \right) [f'(x) - f'(a+b-x)] - \frac{1}{b-a} \int_a^b \left(t - \frac{a+b}{2} \right) f'(t) dt. \end{aligned}$$

Integrating by parts, we have

$$(2.5) \quad \frac{1}{b-a} \int_a^b \left(t - \frac{a+b}{2} \right) f'(t) dt = \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt.$$

Also upon using (2.3), we get

$$(2.6) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b K(t) \left[f'(t) + \left(t - \frac{a+b}{2} \right) f''(t) \right] dt \\ &= \frac{1}{b-a} \int_a^b K(t) f'(t) dt + \frac{1}{b-a} \int_a^b K(t) \left(t - \frac{a+b}{2} \right) f''(t) dt \\ &= \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{b-a} \int_a^b K(t) \left(t - \frac{a+b}{2} \right) f''(t) dt. \end{aligned}$$

It follows from (2.4)–(2.6) that

$$\begin{aligned}
 & \frac{1}{2(b-a)} \int_a^b K(t) \left(t - \frac{a+b}{2} \right) f''(t) dt \\
 &= \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{2} \left[\frac{f(x) + f(a+b-x)}{2} + \frac{f(a) + f(b)}{2} \right] \\
 & \quad + \frac{1}{2} \left(x - \frac{a+b}{2} \right) \frac{f'(x) - f'(a+b-x)}{2}.
 \end{aligned}
 \tag{2.7}$$

Now using (2.7) we obtain

$$\begin{aligned}
 & \left| \frac{1}{2} \left[\frac{f(x) + f(a+b-x)}{2} + \frac{f(a) + f(b)}{2} \right] \right. \\
 & \quad \left. - \frac{1}{2} \left(x - \frac{a+b}{2} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
 & \leq \frac{\|f''\|_\infty}{2(b-a)} \int_a^b |K(t)| \left| t - \frac{a+b}{2} \right| dt.
 \end{aligned}
 \tag{2.8}$$

Since $x \in [a, \frac{a+b}{2}]$, we have

$$\begin{aligned}
 I &:= \int_a^b |K(t)| \left| t - \frac{a+b}{2} \right| dt \\
 &= \int_a^x (t-a) \left| t - \frac{a+b}{2} \right| dt + \int_x^{a+b-x} \left(t - \frac{a+b}{2} \right)^2 dt + \int_{a+b-x}^b (b-t) \left| t - \frac{a+b}{2} \right| dt \\
 &= \int_a^x (t-a) \left(\frac{a+b}{2} - t \right) dt + \int_x^{a+b-x} \left(t - \frac{a+b}{2} \right)^2 dt + \int_{a+b-x}^b (b-t) \left(t - \frac{a+b}{2} \right) dt \\
 &= \frac{(a+3b-4x)(x-a)^2}{12} + \frac{2}{3} \left(\frac{a+b}{2} - x \right)^3 + \frac{(a+3b-4x)(x-a)^2}{12} \\
 &= \frac{(a+3b-4x)(x-a)^2}{6} + \frac{2}{3} \left(\frac{a+b}{2} - x \right)^3,
 \end{aligned}$$

and referring to (2.8), we obtain the result (2.1).

The sharpness of the constant $\frac{1}{3}$ can be proved in a special case for $x = \frac{a+b}{2}$ (see the line behind (1.7)). $\square$

Remark 1. If we take $x = \frac{a+b}{2}$ in (2.1), we recapture the sharp inequality (1.7). If we take $x = a$ in (2.1), we obtain the perturbed trapezoid type inequality

$$\left| \frac{f(a) + f(b)}{2} - \frac{b-a}{8} [f'(b) - f'(a)] - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^2}{24} \|f''\|_\infty,$$

which has a smaller estimator than the sharp trapezoid inequality

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^2}{8} \|f''\|_\infty$$

stated in [11, Proposition 2.7].

Remark 2. Consider

$$F(x) = \left(\frac{a+3b}{4} - x \right) (x-a)^2 + \left(\frac{a+b}{2} - x \right)^3$$

for $x \in [a, \frac{a+b}{2}]$. It's easy to know that $F(x)$ obtains its minimal value at $x = \frac{3a+b}{4}$. Therefore, in (2.1), the point $x = \frac{3a+b}{4}$ gives the best estimator, i.e.,

$$(2.9) \quad \left| \frac{1}{2} \left[\frac{f(\frac{3a+b}{4}) + f(\frac{a+3b}{4})}{2} + \frac{f(a) + f(b)}{2} \right] + \frac{b-a}{8} \frac{f'(\frac{3a+b}{4}) - f'(\frac{a+3b}{4})}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{64} (b-a)^2 \|f''\|_\infty,$$

the right hand side of which is smaller than that of (1.7).

3. APPLICATION TO COMPOSITE QUADRATURE RULES

In [10], the authors utilized inequality (1.6) to give estimates of composite quadrature rules which was pointed out have a markedly smaller error than that which may be obtained by the classical results. In this section, we apply our previous inequality (2.1) to give us estimates of new composite quadrature rules which have a further smaller error.

Theorem 3.1. Let $I_n : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ be a partition of the interval $[a, b]$, $h_i = x_{i+1} - x_i$, $\nu(h) := \max\{h_i : i = 1, \dots, n\}$, $\xi_i \in [x_i, \frac{x_i+x_{i+1}}{2}]$, and

$$S(f, I_n, \xi) = \frac{1}{4} \sum_{i=0}^{n-1} [f(x_i) + f(\xi_i) + f(x_i + x_{i+1} - \xi_i) + f(x_{i+1})] h_i - \frac{1}{4} \sum_{i=0}^{n-1} h_i \left(\xi_i - \frac{x_i + x_{i+1}}{2} \right) [f'(\xi_i) - f'(x_i + x_{i+1} - \xi_i)],$$

then

$$\int_a^b f(x) dx = S(f, I_n, \xi) + R(f, I_n, \xi)$$

and the remainder $R(f, I_n, \xi)$ satisfies the estimate

$$(3.1) \quad |R(f, I_n, \xi)| \leq \frac{1}{3} \|f''\|_\infty \left[\sum_{i=0}^{n-1} \left(\frac{x_i + 3x_{i+1}}{4} - \xi_i \right) (x_i - \xi_i)^2 + \sum_{i=0}^{n-1} \left(\frac{x_i + x_{i+1}}{2} - \xi_i \right)^3 \right].$$

Proof. Inequality (2.1) can be written as

$$(3.2) \quad \left| \int_a^b f(t) dt - \frac{1}{4} [f(a) + f(x) + f(a+b-x) + f(b)] (b-a) + \frac{b-a}{4} \left(x - \frac{a+b}{2} \right) [f'(x) - f'(a+b-x)] \right| \leq \frac{1}{3} \left[\left(\frac{a+3b}{4} - x \right) (x-a)^2 + \left(\frac{a+b}{2} - x \right)^3 \right] \|f''\|_\infty.$$

Applying (3.2) on $\xi_i \in [x_i, \frac{x_i+x_{i+1}}{2}]$, we have

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t) dt - \frac{1}{4} [f(x_i) + f(\xi_i) + f(x_i + x_{i+1} - \xi_i) + f(x_{i+1})] h_i \right. \\ & \quad \left. + \frac{h_i}{4} \left(\xi_i - \frac{x_i + x_{i+1}}{2} \right) [f'(\xi_i) - f'(x_i + x_{i+1} - \xi_i)] \right| \\ & \leq \frac{1}{3} \left[\left(\frac{x_i + 3x_{i+1}}{4} - \xi_i \right) (x_i - \xi_i)^2 + \left(\frac{x_i + x_{i+1}}{2} - \xi_i \right)^3 \right] \|f''\|_\infty. \end{aligned}$$

Now summing over i from 0 to $n-1$ and utilizing the triangle inequality, we have

$$\begin{aligned} & \left| \int_a^b f(t) dt - S(f, I_n, \xi) \right| \\ &= \left| \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} f(t) dt - \frac{1}{4} \sum_{i=0}^{n-1} [f(x_i) + f(\xi_i) + f(x_i + x_{i+1} - \xi_i) + f(x_{i+1})] h_i \right. \\ & \quad \left. + \frac{1}{4} \sum_{i=0}^{n-1} h_i \left(\xi_i - \frac{x_i + x_{i+1}}{2} \right) [f'(\xi_i) - f'(x_i + x_{i+1} - \xi_i)] \right| \\ & \leq \frac{1}{3} \|f''\|_\infty \sum_{i=0}^{n-1} \left[\left(\frac{x_i + 3x_{i+1}}{4} - \xi_i \right) (x_i - \xi_i)^2 + \left(\frac{x_i + x_{i+1}}{2} - \xi_i \right)^3 \right] \end{aligned}$$

and therefore (3.1) holds. $\square$

Corollary 3.1. *If we choose $\xi_i = \frac{3x_i+x_{i+1}}{4}$, then we have*

$$\begin{aligned} \bar{S}(f, I_n) &= \frac{1}{4} \sum_{i=0}^{n-1} \left[f(x_i) + f\left(\frac{3x_i+x_{i+1}}{4}\right) + f\left(\frac{x_i+3x_{i+1}}{4}\right) + f(x_{i+1}) \right] h_i \\ & \quad + \frac{1}{16} \sum_{i=0}^{n-1} \left[f'\left(\frac{3x_i+x_{i+1}}{4}\right) - f'\left(\frac{x_i+3x_{i+1}}{4}\right) \right] h_i^2 \end{aligned}$$

and

$$(3.3) \quad |\bar{R}(f, I_n)| \leq \frac{1}{64} \|f''\|_\infty \sum_{i=0}^{n-1} h_i^3.$$

Remark 3. *It is obvious that inequality (3.3) is better than [10, inequality (3.1)] due to a smaller error.*

4. APPLICATION TO PROBABILITY DENSITY FUNCTIONS

Now, let X be a random variable taking values in the finite interval $[a, b]$, with the probability density function $f : [a, b] \rightarrow [0, 1]$ and with the cumulative distribution function

$$F(x) = \Pr(X \leq x) = \int_a^x f(t) dt.$$

The following result holds:

Theorem 4.1. *With the above assumptions, we have the inequality*

$$(4.1) \quad \left| \frac{1}{2} \left[\frac{F(x) + F(a+b-x)}{2} + \frac{1}{2} \right] - \frac{1}{2} \left(x - \frac{a+b}{2} \right) \frac{f(x) - f(a+b-x)}{2} - \frac{b - E(X)}{b-a} \right| \leq \left[\frac{(\frac{a+3b}{4} - x)(x-a)^2}{3(b-a)^3} + \frac{1}{3} \frac{(\frac{a+b}{2} - x)^3}{(b-a)^3} \right] (b-a)^2 \|f'\|_\infty$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. Follows by (2.1) on choosing $f = F$ and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t) dt,$$

we obtain (4.1). □

In particular, we have:

Corollary 4.1. *With the above assumptions, we have the inequality*

$$\left| \frac{1}{2} \left[\frac{F(\frac{3a+b}{4}) + F(\frac{a+3b}{4})}{2} + \frac{1}{2} \right] + \frac{b-a}{8} \frac{f(\frac{3a+b}{4}) - f(\frac{a+3b}{4})}{2} - \frac{b - E(X)}{b-a} \right| \leq \frac{1}{64} (b-a)^2 \|f'\|_\infty.$$

Acknowledgments. This work was partly supported by the National Natural Science Foundation of China (Grant No. 40975002) and the Natural Science Foundation of the Jiangsu Higher Education Institutions (Grant No. 09KJB110005). The author would like to thank Professor J. Duoandikoetxea for bringing reference [11] to his attention.

REFERENCES

- [1] M. W. Alomari, A companion of Dragomir's generalization of Ostrowski's inequality and applications in numerical integration, RGMIA Res. Rep. Coll., 14 (2011) article 50.
- [2] M. W. Alomari, A companion of Ostrowski's inequality with applications, Transylv. J. Math. Mech. **3** (2011), no. 1, 9–14.
- [3] M. W. Alomari, A companion of Ostrowski's inequality for mappings whose first derivatives are bounded and applications in numerical integration, RGMIA Res. Rep. Coll., 14 (2011) article 57.
- [4] M. W. Alomari, A generalization of companion inequality of Ostrowski's type for mappings whose first derivatives are bounded and applications in numerical integration, RGMIA Res. Rep. Coll., 14 (2011) article 58.
- [5] N. S. Barnett, S. S. Dragomir and I. Gomm, A companion for the Ostrowski and the generalised trapezoid inequalities, Math. Comput. Modelling **50** (2009), no. 1-2, 179–187.
- [6] S. S. Dragomir, A companions of Ostrowski's inequality for functions of bounded variation and applications, RGMIA Res. Rep. Coll., 5 (2002), Supp., article 28.
- [7] S. S. Dragomir, Ostrowski type inequalities for functions defined on linear spaces and applications for semi-inner products, J. Concr. Appl. Math. **3** (2005), no. 1, 91–103.
- [8] S. S. Dragomir, Some companions of Ostrowski's inequality for absolutely continuous functions and applications, Bull. Korean Math. Soc. **42** (2005), no. 2, 213–230.
- [9] S. S. Dragomir, Ostrowski's type inequalities for continuous functions of selfadjoint operators on Hilbert spaces: a survey of recent results, Ann. Funct. Anal. **2** (2011), no. 1, 139–205.
- [10] S. S. Dragomir and A. Sofo, An integral inequality for twice differentiable mappings and applications, Tamkang J. Math. **31** (2000), no. 4, 257–266.
- [11] J. Duoandikoetxea, A unified approach to several inequalities involving functions and derivatives, Czechoslovak Math. J. **51(126)** (2001), no. 2, 363–376.

- [12] A. Guessab and G. Schmeisser, Sharp integral inequalities of the Hermite-Hadamard type, *J. Approx. Theory* **115** (2002), no. 2, 260–288.
- [13] Vu Nhat Huy and Q. -A. Ngô, New bounds for the Ostrowski-like type inequalities, *Bull. Korean Math. Soc.* **48** (2011), no. 1, 95–104.
- [14] Z. Liu, Some companions of an Ostrowski type inequality and applications, *JIPAM. J. Inequal. Pure Appl. Math.* **10** (2009), no. 2, Article 52, 12 pp.
- [15] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and new inequalities in analysis*, Mathematics and its Applications (East European Series), 61, Kluwer Acad. Publ., Dordrecht, 1993.
- [16] M. Z. Sarikaya, On the Ostrowski type integral inequality, *Acta Math. Univ. Comenian. (N.S.)* **79** (2010), no. 1, 129–134.
- [17] M. Z. Sarikaya, New weighted Ostrowski and Čebyšev type inequalities on time scales, *Comput. Math. Appl.* **60** (2010), no. 5, 1510–1514.
- [18] E. Set and M. Z. Sarikaya, On the generalization of Ostrowski and Grüss type discrete inequalities, *Comput. Math. Appl.* **62** (2011), no. 1, 455–461.
- [19] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, Generalizations of weighted Ostrowski type inequalities for mappings of bounded variation and their applications, *Comput. Math. Appl.* **55** (2008), no. 8, 1785–1793.
- [20] K.-L. Tseng, S.-R. Hwang, G.-S. Yang and Y.-M. Chou, Weighted Ostrowski integral inequality for mappings of bounded variation, *Taiwanese J. Math.* **15** (2011), no. 2, 573–585.
- [21] S. W. Vong, A note on some Ostrowski-like type inequalities, *Comput. Math. Appl.* **62** (2011), no. 1, 532–535.

(W. J. Liu) COLLEGE OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, CHINA
E-mail address: wjliu@nuist.edu.cn